

## OBSERVED BEHAVIOR OF HIGHLY INELASTIC ELECTRON-PROTON SCATTERING

M. Breidenbach, J. I. Friedman, and H. W. Kendall

Department of Physics and Laboratory for Nuclear Science,\*  
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

and

E. D. Bloom, D. H. Coward, H. DeStaeblcr, J. Drees, L. W. Mo, and R. E. Taylor

Stanford Linear Accelerator Center,† Stanford, California 94305

(Received 22 August 1969)

Results of electron-proton inelastic scattering at  $6^\circ$  and  $10^\circ$  are discussed, and values of the structure function  $W_2$  are estimated. If the interaction is dominated by transverse virtual photons,  $\nu W_2$  can be expressed as a function of  $\omega = 2M\nu/q^2$  within experimental errors for  $q^2 > 1$  (GeV/c) $^2$  and  $\omega > 4$ , where  $\nu$  is the invariant energy transfer and  $q^2$  is the invariant momentum transfer of the electron. Various theoretical models and sum rules are briefly discussed.

In a previous Letter,<sup>1</sup> we have reported experimental results from a Stanford Linear Accelerator Center-Massachusetts Institute of Technology study of high-energy inelastic electron-proton scattering. Measurements of inelastic spectra, in which only the scattered electrons were detected, were made at scattering angles of  $6^\circ$  and  $10^\circ$  and with incident energies between 7 and 17 GeV. In this communication, we discuss some of the salient features of inelastic spectra in the deep continuum region.

One of the interesting features of the measurements is the weak momentum-transfer dependence of the inelastic cross sections for excitations well beyond the resonance region. This weak dependence is illustrated in Fig. 1. Here we have plotted the differential cross section divided by the Mott cross section,  $(d^2\sigma/d\Omega dE')/(\sigma_{\text{Mott}})$ , as a function of the square of the four-momentum transfer,  $q^2 = 2EE'(1 - \cos\theta)$ , for constant values of the invariant mass of the recoiling target system,  $W$ , where  $W^2 = 2M(E - E') + M^2 - q^2$ .  $E$  is the energy of the incident electron,  $E'$  is the energy of the final electron, and  $\theta$  is the scattering angle, all defined in the laboratory system;  $M$  is the mass of the proton. The cross section is divided by the Mott cross section

$$\left(\frac{d\sigma}{d\Omega}\right)_{\text{Mott}} = \frac{e^4}{4E^2} \frac{\cos^2 \frac{1}{2}\theta}{\sin^4 \frac{1}{2}\theta}$$

in order to remove the major part of the well-known four-momentum transfer dependence arising from the photon propagator. Results from both  $6^\circ$  and  $10^\circ$  are included in the figure for each value of  $W$ . As  $W$  increases, the  $q^2$  dependence appears to decrease. The striking difference

between the behavior of the inelastic and elastic cross sections is also illustrated in Fig. 1, where the elastic cross section, divided by the Mott cross section for  $\theta = 10^\circ$ , is included. The  $q^2$  dependence of the deep continuum is also consider-




FIG. 1.  $(d^2\sigma/d\Omega dE')/\sigma_{\text{Mott}}$ , in  $\text{GeV}^{-1}$ , vs  $q^2$  for  $W = 2, 3$ , and  $3.5$  GeV. The lines drawn through the data are meant to guide the eye. Also shown is the cross section for elastic  $e$ - $p$  scattering divided by  $\sigma_{\text{Mott}}$ ,  $(d\sigma/d\Omega)/\sigma_{\text{Mott}}$ , calculated for  $\theta = 10^\circ$ , using the dipole form factor. The relatively slow variation with  $q^2$  of the inelastic cross section compared with the elastic cross section is clearly shown.

ably weaker than that of the electroexcitation of the resonances,<sup>2</sup> which have a  $q^2$  dependence similar to that of elastic scattering for  $q^2 > 1$  (GeV/c)<sup>2</sup>.

On the basis of general considerations, the differential cross section for inelastic electron scattering in which only the electron is detected can be represented by the following expression<sup>3</sup>:

$$\frac{d^2\sigma}{d\Omega dE'} = \left(\frac{d\sigma}{d\Omega}\right)_{\text{Mott}} (W_2 + 2W_1 \tan^2 \frac{1}{2}\theta).$$

The form factors  $W_2$  and  $W_1$  depend on the properties of the target system, and can be represented as functions of  $q^2$  and  $\nu = E - E'$ , the electron energy loss. The ratio  $W_2/W_1$  is given by

$$\frac{W_2}{W_1} = \left(\frac{q^2}{\nu^2 + q^2}\right) (1 + R), \quad R \geq 0,$$

where  $R$  is the ratio of the photoabsorption cross sections of longitudinal and transverse virtual photons,  $R = \sigma_S/\sigma_T$ .<sup>4</sup>

The objective of our investigations is to study the behavior of  $W_1$  and  $W_2$  to obtain information about the structure of the proton and its electromagnetic interactions at high energies. Since at present only cross-section measurements at small angles are available, we are unable to

make separate determinations of  $W_2$  and  $W_1$ . However, we can place limits on  $W_2$  and study the behavior of these limits as a function of the invariants  $\nu$  and  $q^2$ .

Bjorken<sup>5</sup> originally suggested that  $W_2$  could have the form

$$W_2 = (1/\nu)F(\omega),$$

where

$$\omega = 2M\nu/q^2.$$

$F(\omega)$  is a universal function that is conjectured to be valid for large values of  $\nu$  and  $q^2$ . This function is universal in the sense that it manifests scale invariance, that is, it depends only on the ratio  $\nu/q^2$ . Since

$$\nu W_2 = \frac{\nu d^2\sigma/d\Omega dE'}{(d\sigma/d\Omega)_{\text{Mott}}} \left[1 + 2 \frac{1}{1+R} \left(1 + \frac{\nu^2}{q^2}\right) \tan^2 \frac{1}{2}\theta\right]^{-1},$$

the value of  $\nu W_2$  for any given measurement clearly depends on the presently unknown value of  $R$ . It should be noted that the sensitivity to  $R$  is small when  $2(1 + \nu^2/q^2) \tan^2 \frac{1}{2}\theta \ll 1$ . Experimental limits on  $\nu W_2$  can be calculated on the basis of the extreme assumptions  $R=0$  and  $R=\infty$ . In Figs. 2(a) and 2(b) the experimental values of  $\nu W_2$




FIG. 2.  $\nu W_2$  vs  $\omega = 2M\nu/q^2$  is shown for various assumptions about  $R = \sigma_S/\sigma_T$ . (a) 6° data except for 7-GeV spectrum for  $R=0$ . (b) 10° data for  $R=0$ . (c) 6° data except for 7-GeV spectrum for  $R=\infty$ . (d) 10° data for  $R=\infty$ . (e) 6°, 7-GeV spectrum for  $R=0$  and  $R=\infty$ .

from the  $6^\circ$  and  $10^\circ$  data for  $q^2 > 0.5$  (GeV/c) $^2$  are shown as a function of  $\omega$  for the assumption that  $R = 0$ . Figures 2(c) and 2(d) show the experimental values of  $\nu W_2$  calculated from the  $6^\circ$  and  $10^\circ$  data with  $q^2 > 0.5$  (GeV/c) $^2$  under the assumption  $R = \infty$ . The  $6^\circ$ , 7-GeV results for  $\nu W_2$ , all of which have values of  $q^2 \leq 0.5$  (GeV/c) $^2$ , are shown for both assumptions in Fig. 2(e). The elastic peaks are not displayed in Fig. 2.

The results shown in these figures indicate the following:

(1) If  $\sigma_T \gg \sigma_S$ , the experimental results are consistent with a universal curve for  $\omega \gtrsim 4$  and  $q^2 \gtrsim 0.5$  (GeV/c) $^2$ . Above these values, the measurements at  $6^\circ$  and  $10^\circ$  give the same results within the errors of measurements. The  $6^\circ$ , 7-GeV measurements of  $\nu W_2$ , all of which have values of  $q^2 \leq 0.5$  (GeV/c) $^2$ , are somewhat smaller than the results from the other spectra in the continuum region.

The values of  $\nu W_2$  for  $\omega \gtrsim 5$  show a gradual decrease as  $\omega$  increases. In order to test the statistical significance of the observed slope, we have made linear least-squares fits to the values of  $\nu W_2$  in the region  $6 \leq \omega \leq 25$ . These fits give  $\nu W_2 = (0.351 \pm 0.023) - (0.00386 \pm 0.00088)\omega$  for data with  $q^2 > 0.5$  (GeV/c) $^2$  and  $\nu W_2 = (0.366 \pm 0.024) - (0.0045 \pm 0.0019)\omega$  for  $q^2 > 1$  (GeV/c) $^2$ . The quoted errors consist of the errors from the fit added in quadrature with estimates of systematic errors.

Since  $\sigma_T + \sigma_S \simeq 4\pi^2\alpha\nu W_2/q^2$  for  $\omega \gg 1$ , our results can provide information about the behavior of  $\sigma_T$  if  $\sigma_T \gg \sigma_S$ . The scale invariance found in the measurements of  $\nu W_2$  indicates that the  $q^2$  dependence of  $\sigma_T$  is approximately  $1/q^2$ . The gradual decrease exhibited in  $\nu W_2$  for large  $\omega$  suggests that the photoabsorption cross section for virtual photons falls slowly at constant  $q^2$  as the photon energy  $\nu$  increases.

The measurements indicate that  $\nu W_2$  has a broad maximum in the neighborhood of  $\omega = 5$ . The question of whether this maximum has any correspondence to a possible quasielastic peak<sup>6</sup> requires further investigation.

It should be emphasized that all of the above conclusions are based on the assumption that  $\sigma_T \gg \sigma_S$ .

(2) If  $\sigma_S \gg \sigma_T$ , the measurements of  $\nu W_2$  do not follow a universal curve and have the general feature that at constant  $2M\nu/q^2$ , the value of  $\nu W_2$  increases with  $q^2$ .

(3) For either assumption,  $\nu W_2$  shows a threshold behavior in the range  $1 \leq \omega \leq 4$ .  $W_2$  is con-

strained to be zero at inelastic threshold which corresponds to  $\omega \simeq 1$  for large  $q^2$ . In the threshold region of  $\nu W_2$ ,  $W_2$  falls rapidly as  $q^2$  increases at constant  $\nu$ . This qualitatively different from the weak  $q^2$  behavior for  $\omega > 4$ . For  $q^2 \approx 1$  (GeV/c) $^2$ , the threshold region contains the resonances excited in electroproduction. As  $q^2$  increases, the variations due to these resonances damp out and the values of  $\nu W_2$  do not appear to vary rapidly with  $q^2$  at constant  $\omega$ .

It can be seen from a comparison of Figs. 2(a) and 2(c) that the  $6^\circ$  data provide a measurement of  $\nu W_2$  to within 10% up to a value of  $\omega \approx 6$ , irrespective of the values of  $R$ .

There have been a number of different theoretical approaches in the interpretation of the high-energy inelastic electron-scattering results. One class of models,<sup>6-9</sup> referred to as parton models, describes the electron as scattering incoherently from pointlike constituents within the proton. Such models lead to a universal form for  $\nu W_2$ , and the point charges assumed in specific models give the magnitude of  $\nu W_2$  for  $\omega > 2$  to within a factor of 2.<sup>6</sup> Another approach<sup>10,11</sup> relates the inelastic scattering to off-the-mass-shell Compton scattering which is described in terms of Regge exchange using the Pomeranchuk trajectory. Such models lead to a flat behavior of  $\nu W_2$  as a function of  $\nu$  but do not require the weak  $q^2$  dependence observed and do not make any numerical predictions at this time. Perhaps the most detailed predictions made at present come from a vector-dominance model which primarily utilizes the  $\rho$  meson.<sup>12</sup> This model reproduces the gross behavior of the data and has the feature that  $\nu W_2$  asymptotically approaches a function of  $\omega$  as  $q^2 \rightarrow \infty$ . However, a comparison of this model with the data leads to statistically significant discrepancies. This can be seen by noting that the prediction for  $d^2\sigma/d\Omega dE'$  contains a parameter  $\xi$ , the ratio of the cross sections for longitudinally and transversely polarized  $\rho$  mesons on protons, which is expected to be a function of  $W$  but which should be independent of  $q^2$ . For values of  $W \geq 2$  GeV, the experimental values of  $\xi$  increase by about  $(50 \pm 5)\%$  as  $q^2$  increases from 1 to 4 (GeV/c) $^2$ . This model predicts that

$$\sigma_S/\sigma_T = \xi(W)(q^2/m_\rho^2)[1 - q^2/2m\nu],$$

which will provide the most stringent test of this approach when a separation of  $W_1$  and  $W_2$  can be made.

The application of current algebra<sup>13-17</sup> and the use of current commutators leading to sum rules

and sum-rule inequalities provide another way of comparing the measurements with theory. There have been some recent theoretical considerations<sup>18-20</sup> which have pointed to possible ambiguity in these calculations; however, it is still of considerable interest to compare them with experiment.

In general,  $W_2$  and  $W_1$  can be related to commutators of electromagnetic current densities.<sup>6,16</sup> The experimental value of the energy-weighted sum  $\int_1^\infty (d\omega/\omega^2)(\nu W_2)$ , which is related to the equal-time commutator of the current and its time derivative, is  $0.16 \pm 0.01$  for  $R=0$  and  $0.20 \pm 0.03$  for  $R=\infty$ . The integral has been evaluated with an upper limit  $\omega=20$ . This integral is also important in parton theories where its value is the mean square charge per parton.

Gottfried<sup>21</sup> has calculated a constant- $q^2$  sum rule for inelastic electron-proton scattering based on a nonrelativistic quark model involving pointlike quarks. The resulting sum rule is

$$\int_1^\infty \frac{d\omega}{\omega} (\nu W_2) = \int_{q^2/2M}^\infty d\nu W_2$$

$$= 1 - \frac{G_{Ep}^2 + (q^2/4M^2)G_{Mp}}{1 + q^2/4M^2},$$

where  $G_{Ep}$  and  $G_{Mp}$  are the electric and magnetic form factors of the proton. The experimental evaluation of this integral from our data is much more dependent on the assumption about  $R$  than the previous integral. We will thus use the 6° measurements of  $W_2$  which are relatively insensitive to  $R$ . Our data for a value of  $q^2 \approx 1$  (GeV/c)<sup>2</sup>, which extend to a value of  $\nu$  of about 10 GeV, give a sum that is  $0.72 \pm 0.05$  with the assumption that  $R=0$ . For  $R=\infty$ , its value is  $0.81 \pm 0.06$ . An extrapolation of our measurements of  $\nu W_2$  for each assumption suggests that the sum is saturated in the region  $\nu \approx 20$ -40 GeV. Bjorken<sup>13</sup> has proposed a constant- $q^2$  sum-rule inequality for high-energy scattering from the proton and neutron derived on the basis of current algebra. His result states that

$$\int_1^\infty \frac{d\omega}{\omega} \nu (W_{2p} + W_{2n}) = \int_{q^2/2M}^\infty d\nu (W_{2p} + W_{2n}) \geq \frac{1}{2},$$

where the subscripts  $p$  and  $n$  refer to the proton and neutron, respectively. Since there are presently no electron-neutron inelastic scattering results available, we estimate  $W_{2n}$  in a model-dependent way. For a quark model<sup>22</sup> of the proton,  $W_{2n} \approx 0.8W_{2p}$  whereas in the model<sup>8</sup> of Drell and co-workers,  $W_{2n}$  rapidly approaches  $W_{2p}$  as  $\nu$  in-

creases. Using our results, this inequality is just satisfied at  $\omega \approx 4.5$  for the quark model and at  $\omega \approx 4.0$  for the other model for either assumption about  $R$ . For example, this corresponds to a value of  $\nu \approx 4.5$  GeV for  $q^2 = 2$  (GeV/c)<sup>2</sup>. Bjorken<sup>23</sup> estimates that the experimental value of the sum is too small by about a factor of 2 for either model, but it should be noted that the  $q^2$  dependence found in the data is consistent with the predictions of this calculation.

---

\*Work supported in part through funds provided by the U. S. Atomic Energy Commission under Contract No. AT(30-1)2098.

†Work supported by the U. S. Atomic Energy Commission.

<sup>1</sup>E. Bloom et al., preceding Letter [Phys. Rev. Letters **23**, 930 (1969)].

<sup>2</sup>Preliminary results from the present experimental program are given in the report by W. K. H. Panofsky, in Proceedings of the Fourteenth International Conference On High Energy Physics, Vienna, Austria, 1968 (CERN Scientific Information Service, Geneva, Switzerland, 1968), p. 23.

<sup>3</sup>R. von Gehlen, Phys. Rev. **118**, 1455 (1960); J. D. Bjorken, 1960 (unpublished); M. Gourdin, Nuovo Cimento **21**, 1094 (1961).

<sup>4</sup>See L. Hand, in Proceedings of the Third International Symposium on Electron and Photon Interactions at High Energies, Stanford Linear Accelerator Center, Stanford, California, 1967 (Clearing House of Federal Scientific and Technical Information, Washington, D. C., 1968), or F. J. Gilman, Phys. Rev. **167**, 1365 (1968).

<sup>5</sup>J. D. Bjorken, Phys. Rev. **179**, 1547 (1969).

<sup>6</sup>J. D. Bjorken and E. A. Paschos, Stanford Linear Accelerator Center, Report No. SLAC-PUB-572, 1969 (to be published).

<sup>7</sup>R. P. Feynman, private communication.

<sup>8</sup>S. J. Drell, D. J. Levy, and T. M. Yan, Phys. Rev. Letters **22**, 744 (1969).

<sup>9</sup>K. Huang, in Argonne National Laboratory Report No. ANL-HEP 6909, 1968 (unpublished), p. 150.

<sup>10</sup>H. D. Abarbanel and M. L. Goldberger, Phys. Rev. Letters **22**, 500 (1969).

<sup>11</sup>H. Harari, Phys. Rev. Letters **22**, 1078 (1969).

<sup>12</sup>J. J. Sakurai, Phys. Rev. Letters **22**, 981 (1969).

<sup>13</sup>J. D. Bjorken, Phys. Rev. Letters **16**, 408 (1966).

<sup>14</sup>J. D. Bjorken, in Selected Topics in Particle Physics, Proceedings of the International School of Physics "Enrico Fermi," Course XLI, edited by J. Steinberger (Academic Press, Inc., New York, 1968).

<sup>15</sup>J. M. Cornwall and R. E. Norton, Phys. Rev. **177**, 2584 (1969).

<sup>16</sup>C. G. Callan, Jr., and D. J. Gross, Phys. Rev. Letters **21**, 311 (1968).

<sup>17</sup>C. G. Callan, Jr., and D. J. Gross, Phys. Rev. Letters **22**, 156 (1969).

<sup>18</sup>R. Jackiw and G. Preparata, Phys. Rev. Letters 22, 975 (1969).

<sup>19</sup>S. L. Adler and W.-K. Tung, Phys. Rev. Letters 22, 978 (1969).

<sup>20</sup>H. Cheng and T. T. Wu, Phys. Rev. Letters 22, 1409

(1969).

<sup>21</sup>K. Gottfried, Phys. Rev. Letters 18, 1174 (1967).

<sup>22</sup>J. D. Bjorken, Stanford Linear Accelerator Center, Report No. SLAC-PUB-571, 1969 (unpublished).

<sup>23</sup>J. D. Bjorken, private communication.

## INCOHERENT PRODUCTION OF $\rho^0$ MESONS FROM NUCLEI AND VECTOR DOMINANCE

G. von Bochmann\* and B. Margolis\*

Deutsches Elektronen-Synchrotron, Hamburg, Germany, and McGill University, Montreal 110, Canada

(Received 22 August 1969)

Incoherent photoproduction of  $\rho^0$  mesons on the nucleus is calculated assuming vector dominance. The results are sensitive to the  $\rho$ -nucleon total cross section which is found to decrease with energy. One finds a relatively weak dependence on energy of the incoherent effective nucleon number.

Recently<sup>1,2</sup> there have been measurements of the incoherent photoproduction of  $\rho^0$  mesons on atomic nuclei at incident photon energies of 2.7, 4, and 8 GeV and momentum transfers  $\sqrt{-t} \sim 0.25$  to 0.35 GeV/c. The relative constancy of the measurements at 4 and 8 GeV is pointed out by the authors of Ref. 2. This could imply some question as to the validity of the vector dominance hypothesis or at least for the eikonal models<sup>3,4</sup> normally used to calculate incoherent production.

The purpose of this note is to point out that (1) there is good reason to believe that the total  $\rho^0$ -nucleon cross section,  $\sigma_\rho$ , decreases with energy, following the  $\pi$ -nucleon cross section closely in value; and (2) as a result of  $\sigma_\rho$  decreasing, and being smaller in value than some authors have deduced, the cross section for incoherent production of  $\rho^0$  mesons in atomic nuclei varies considerably less with energy than one might expect.

Gottfried and Yennie<sup>3</sup> describe incoherent  $\rho^0$  production in terms of a superposition of one- and two-step processes. In the one-step process the  $\rho^0$  meson is produced incoherently on a nucleon and then proceeds, with some damping, to be emitted from the nucleus which has been excited. The two-step process consists of coherent production on one nucleon (no nuclear excitation) followed by incoherent scattering (nuclear excitation occurs) of the  $\rho^0$  meson on another nucleon. The cross section for these processes is given by

$$d\sigma^{(I)}/dt = \int d^2b dz I(b, z) \equiv [d\sigma_0(t)/dt] N_{\text{eff}}, \quad (1)$$

$$I(b, z) = D(b, z) \frac{d\sigma_0(t)}{dt} \exp[-\sigma_\rho \int_z^\infty D(b, z') dz'] \quad (2)$$

$$\varphi(b, z) = -\frac{1}{2}\sigma_\rho \int_{-\infty}^z dz' D(b, z') \exp[-\frac{1}{2}\sigma_\rho \int_{z'}^z D(b, z'') dz''], \quad (3)$$

where  $d\sigma_0(t)/dt$  is the  $\rho^0$  photoproduction differential cross section on a neutron or proton (taken equal) and  $D(b, z)$  is the  $A$ -particle nucleon density function. We neglect the real part of the  $\rho$ -nucleon forward-scattering amplitude. The wave numbers for the photon and  $\rho^0$  meson are  $k_\gamma$  and  $k_\rho$ , respectively. Nuclear c.m. motion and nuclear correlations are neglected. The effective nucleon number  $N_{\text{eff}}$  is a function of  $A$ , energy, and  $\sigma_\rho$ .

At low enough energies ( $\sim 2$  GeV) the one-step process dominates. The two-step process is inhibited due to mismatch of the photon and  $\rho^0$ -meson wave numbers because of the mass of the  $\rho^0$  meson. One has then an incoherent production

cross section

$$\frac{d\sigma^{(I)}(t)}{dt} \simeq \frac{d\sigma_0(t)}{dt} N(A; 0, \sigma_\rho). \quad (4)$$

The effective nucleon numbers  $N(A; \sigma_1, \sigma_2)$  are defined by Kölbig and Margolis.<sup>5</sup> At very high energy where the mass of the  $\rho^0$  is negligible, one finds

$$\frac{d\sigma^{(I)}(t)}{dt} = \frac{d\sigma_0(t)}{dt} N(A; \sigma_\rho, \sigma_\rho). \quad (5)$$

The photon in this case behaves as though it were a  $\rho$  meson. Since  $N(A; \sigma_\rho, \sigma_\rho)$  is considerably less than  $N(A; 0, \sigma_\rho)$  the cross section has fallen in go-